



Short Communication

Effect of Dietary Vitamin A Supplement on Serum Protein of Rats Infected with *Trypanosoma brucei*

Edoga C.O.¹, Njoku O.O.², Ufele A.N.^{1*} and Ebenebe C.I.¹

¹Nnamdi Azikiwe University Awka, Anambra State, NIGERIA

²Federal University of Science and Technology Owerri, Imo State, NIGERIA

Available online at: www.isca.in

Received 25th April 2012, revised 10th August 2012, accepted 15th August 2012

Abstract

The role of dietary vitamin A supplement on the serum protein of trypanosome-infected rats was studied. The rats were inoculated with trypanosomes intraperitoneally and samples were collected on fourth, eighth, twelfth and sixteenth days of post infection. The experiment was carried out at the Department of Biochemistry, Nnamdi Azikiwe University Awka. 60 parasite free-albino rats were used, which were divided into four groups. Group A (control) was left uninfected with trypanosomes, group B and C were infected with trypanosomes and treated with 50mls and 100mls of vitamin A per kg of feed respectively and group D was infected and left untreated with vitamin A. Analyses of the sera using Bradford method and cellulose acetate electrophoresis showed that vitamin A influenced the state of hypoproteinaemia in the trypanosome-infected rats. This was manifested by a positive increase in the level of total serum protein concentration, albumin and beta-globulin. Vitamin A also delayed the proliferation of the parasites associated with trypanosomiasis.

Keywords: Vitamin A, *Trypanosoma brucei*, *Rattus norvegicus*.

Introduction

For several decades, trypanosomiasis has continued to contribute adversely to the economic well being of sub Sahara Africa. Trypanosomiasis is one of the most important livestock diseases in sub-Saharan Africa¹. This scourge remains a pressing challenge especially to African scientists in formulating a possible action plan that would be basic on the poor resources of the African continent. The articulation of such plan would include both preventive measures and treatment modalities

Trypanosome is known to attack red blood cells and vascular endothelium. It concentrates more in the peripheral circulation². These trypanosomes exhibit remarkable antigenic variation of their surface glycoprotein with hundreds of antigenic type. One antigenic type will coat the surface of the parasites for approximately 10 days followed by other types in sequences in the new progeny. This variation is due to sequential movement of the glycoprotein genes to a preferential location of the chromosome were only that specific gene is transcribed into messenger RNA. These antigenic variations allow the organism to continually invade the host immune response. The first wave of parasitemia is accompanied by depressed packed cell volume, neutropenia and thrombocytopenia³.

Trypanosomiasis in rats is associated with a decreased serum protein as infection progressed. Improvement on host's nutrition is important in moderating the severity of pathophysiological effect of trypanosomiasis and also influences the rate of recovery⁴. It was also discovered that

supplementary feeding significantly reduces the severity of trypanosomiasis^{5,6}. Vitamin A is essential for the development of bursa of fabricius, thymus and immunity in chickens⁷. Vitamin A supplements strongly increase serum protein and glycoprotein synthesis which maintain synthesis of surface coating mucopolysaccharides. Hence increase mucosal immune system, together with the epithelial rebuilding character of vitamin A raise macrophage activity and increase humoral and cell mediated response of vitamin A sufficient chicken, strongly suggest a much high resistance to infection of higher animals^{8,9}.

Over the years, vitamin A has been used to tackle all kinds of infections and it is very popular both among the low, middle and higher socio-economic class^{10,11}. This study was design to evaluate to what extent this vitamin A can influence the state of hypoproteinaemia in trypanosome-infected rats.

Material and Methods

Twenty 90-day old male albino rats (*Rattus norvegicus*) weighing approximately 145g, were used for this experiment. The rats were marked for identification and held in stainless wire-rats-cages in clean experimental animal house. The cages were labeled A to D corresponding to four groups and each group had five rats. Diet 1 was given to rats in Cage A which contained 1kg of chick mash without vitamin A. Diet 2 was given to rats in Cage B which contained 1kg of chick mash mixed with 50mls of vitamin A. Diet 3 was used to feed rats in Cage C which contained 1kg of chick mash mixed with 100mls of vitamin A and Diet 4 was used to feed rats in Cage D which

contained 1kg of chick mash without any vitamin A. Rats in Cage A were not infected while rats in Cages B, C and D were infected with *Trypanosoma brucei*. One rat was first inoculated with trypanosome of NITR type from Veterinary Medicine Faculty, University of Nigeria, Nsukka. It was isolated from other animals and after 14 days of inoculation, the blood of that rat was used to inoculate others. Each experimental rat that was inoculated was given 0.1ml of infected blood in normal saline, which contained about eight thousand trypanosomes, using a matching chart¹² to determine the level of parasitemia. Rats in Cages A and D served as control groups. Each experimental set up was replicated three times. The rats had unlimited supply of clean water.

5ml of the blood of the rats were collected in each experimental day which was four days intervals for sixteen days of the experimental period to determine the total serum protein. The collected blood was allowed to clot for about 30 minutes at room temperature. Then each sample was centrifuged at 3,000rpm for 10 minutes and the serum was removed. The sera were used immediately for serum protein determination using Bradford method. The absorbance of the solutions was read at 520nm-wavelengths using spectrophotometer.

Statistical analysis: The change in total serum protein level was determined using 2-way ANOVA and data obtained were reported as mean concentration.

Results and Discussion

The result obtained indicated that administration of vitamin A positively influenced the serum protein of trypanosome-infected rats. There was increase in the total serum protein, raised albumin and beta-globulin in the infected and treated rats. This was shown in tables 1, 2 and 3. Table - 1 shows the level of total serum concentration of the experimental rats. Table - 2 shows the albumin (g/l) of the experimental rats while table - 3 shows the Beta-globulin (g/l) of the experimental rats. The lowest level of total serum protein of 49.27g/l, 17.85g/l albumin and 3.46g/l beta-globulin was observed in infected and untreated rats, which was Cage D. Followed by 50.79g/l total serum protein, 20.77g/l albumin and 5.15g/l beta-globulin observed in infected rats treated with 50ml of vitamin A per kg of chick mash, which was Cage B. Then infected rats which were treated with 100ml of vitamin A per kg of chick mash had 52.2g/l of total serum protein, 22.99g/l albumin and 5.36g/l beta-globulin which was Cage C. The highest level was seen in rats in Cage A which had 60.99g/l total serum protein, 32.24g/l albumin and 8.26g/l beta-globulin.

Several scientific researches have been done on trying to identify and standardize active food supplement that would be active in treatment of trypanosomiasis. This trypanosomiasis has contributed adversely to the economic and social well being of Sub-Sahara Africans. A lot of scientific investigations have proved that retinol and retinoic acid nutrient produced significant enhancement on the immune system of rats¹³.

Table-1
Total serum (g/l) of the experimental rats with the length of post-infection

Length of post-infection	Group A (uninfected and untreated)	Group B (infected and treated with 50ml vitamin A/kg of feed)	Group C (infected and treated with 100ml vitamin A/kg of feed)	Group D (infected and untreated)
4	60.63	59.80	59.82	59.53
8	59.84	55.88	56.24	54.04
12	61.65	50.83	52.46	48.79
16	61.86	36.67	40.31	34.71
Total	243.98	203.18	208.83	199.07
Mean	60.99	50.79	52.21	49.27

Table-2
Albumin (g/l) of the experimental rats with the length of post-infection

Length of post-infection	Group A (uninfected and untreated)	Group B (infected and treated with 50ml vitamin A/kg of feed)	Group C (infected and treated with 100ml vitamin A/kg of feed)	Group D (infected and untreated)
4	32.68	30.02	30.21	27.40
8	30.93	20.43	24.62	16.51
12	32.69	19.75	21.06	15.73
14	32.64	12.87	16.09	11.74
Total	128.94	83.07	91.98	71.38
Mean	32.24	20.77	22.99	17.85

Table-3
Beta-globulin (g/l) of the experimental rats with the length of post-infection

Length of post-infection	Group A (uninfected and untreated)	Group B (infected and treated with 50ml vitamin A/kg of feed)	Group C (infected and treated with 100ml vitamin A/kg of feed)	Group D (infected and untreated)
4	7.88	6.45	6.92	5.84
8	8.66	5.18	5.77	3.19
12	8.16	4.21	4.86	2.68
14	8.35	4.76	3.91	2.12
Total	33.05	20.60	21.46	13.83
Mean	8.26	5.15	5.36	3.46

The observed effect of vitamin A supplement from this study on the serum protein of trypanosome-infected rats is attributed to its effect on the haemopoietic system. The effect of vitamin A on the infected and treated rats when compared with the infected and untreated rats showed that vitamin A had positive influence on the defense capacity of infected and treated animals. Therefore, this study has provided evidence that vitamin A has a potential for influencing the state of hypoproteinaemia in the trypanosome-infected rats. Even if vitamin A cannot destroy the trypanosomes, it can ameliorate the stress of trypanosomiasis and as enhanced the hosts' immune system to fight the invaded pathogens.

Conclusion

From the research, it is advisable to include vitamin A in the feed of livestock because it enhances the immune system of animal. Also in the treatment of trypanosomiasis, it is advisable to give vitamin A to infected animals for quick recovery.

References

- Morrison W.I., Murray M. and McIntyre W.I.M., Bovine trypanosomiasis, In Risstic M. and McIntyre, W.I.M. (Eds), *Disease of Cattle in Tropics*, The Hague, Martinis Nijhoff Publishers, 469-497 (1981)
- Jackson G.J., Trypanosoma congolense: inheritance of susceptibility to infection in inbred strains of mice, *Experimental Parasitology*, **48**, 378-383 (1979)
- Krampitz H.E., Beobach, tungen an experimentallan infection ostafrikkascher Zebundider Mit wild Stamen Von, T. congolense, *Trop. Para.*, **21**, 1-30 (1970)
- Katungka-Rwakishaya E., Interaction between animal nutrition and parasites, studies with experimental trypanosomiasis in sheep, Pages 1-9, In (Lebbie, S.H. B. and Kagwini, E., Eds), *Small Ruminant Research and Development in Africa*, International Livestock Research Institute (ILRI) Nairobi Kenya (1996)
- Agyemang K., Dwinger R.H., Touray B.N., Jeannin P., Fofana D. and Grieve A.S., Effects of Nutrition on degree of aneamia and live weight changes in N'Dama cattle infected with trypanosomes, *Livestock Production Science*, **26**, 39-51 (1990)
- Little D.A., Dwinger R.H., Clifford D.J., Grieve A.S., Kora S. and Bojang M., Effect of Nutritional level and body condition on susceptibility of N'Dama cattle to T. congolense infection in the Gambia, *Proceedings of the Nutrition society*, **49**, 209A (1990)
- Raza A., Khan F.K., Saced M.A. and Basir N.I., Effect of vitamin A on Growth Traits, Immuno-regulatory Organs and Immune Response in Broiler Chicken, *J. Applied Ani. Res.*, **12**, 81-88 (1997)
- Sijtsma R.S., West F.C., Pombout M.W.J. and Van Der Zipp J.A., The interaction between vitamin A status and Newcastle Disease virus infections in chicken, *J. Nutr.*, **119**, 932-939 (1991)
- Johannsen A.K.B., Jensen S.K. and Jakobsen K., A note on vitamin A activity of B-Carotene in rats, *Acta Agric. Scand. Sect. J. Ani. Sc.*, **48**, 260-263 (1998)
- Bang F.B., Bang B.G. and Foard M., Acute Newcastle viral infection of the upper respiratory tract of the chicken: 2, Effect of diets deficient in vitamin A on the infection, *J. Ani. Path.*, **78**, 417-426 (1995)
- Benynen A.C., Sijtsma R.S., Kiepurski A.K., West C.E., Baumans V., Herck Van. H., Stafleu R.F. and Van Tintele G., Objective clinical examination of poultry as illustrated by the comparison of chicken with different vitamin A status, *British J. An. Sc.*, **23**, 307-312 (1989)
- Herbert W.J. and Lumbsden W.H.R., Trypanosoma brucei, A rapid matching method for estimating the host's parasitemia, *Expt. Para.*, **40**, 427-431 (1976)
- Nauss K.M., Phua C.C., Ambrodi L. and Newberne P.M., Immunological changes during progressive stages of vitamin A deficiency in the rats, *J. Nutr.*, **115**, 909-918 (1985)